

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 September 2001 (27.09.2001)

PCT


(10) International Publication Number
WO 01/71012 A1

- (51) International Patent Classification⁷: C12N 15/77. 15/53, 9/02, C12P 13/08
- (21) International Application Number: PCT/EP00/06299
- (22) International Filing Date: 5 July 2000 (05.07.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/531.265 20 March 2000 (20.03.2000) US
- (71) Applicants: DEGUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE). NATIONAL UNIVERSITY OF IRELAND [IE/IE]; Galway (IE).
- (72) Inventors: DUNICAU, L., K. (deceased). MCCORMACK, Ashling; Moate Road, Athlone, County Westmeath (IE). STAPELTON, Cliona; 27 Railway View, Roscrea, County Tipperary (IE). BURKE, Kevin; 5 Greenfield Road, Newcastle, Galway, County Galway (IE). MÖCKEL, Bettina; Benrodestrasse 35, D-40597 Düsseldorf (DE).
- (81) Designated States (*national*): AU, BR, CA, CN, HU, ID, JP, KR, MX, PL, RU, SK, UA, ZA.
- (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: PROCESS FOR THE FERMENTATIVE PREPARATION OF L-AMINO ACIDS WITH AMPLIFICATION OF THE GND GENE

Map of the plasmid pECgnd



(57) Abstract: The invention relates to a process for the preparation of L-amino acids by fermentation of coryneform bacteria, which comprises carrying out the following steps: a) fermentation of the desired L-amino acid-producing bacteria in which at least the gnd gene is amplified, b) concentration of the L-amino acid in the medium or in the cells of the bacteria and c) isolation of the L-amino acid produced.

WO 01/71012 A1



— with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**Process for the fermentative preparation of L-amino acids
with amplification of the gnd gene**

The invention relates to a process for the fermentative preparation of L-amino acids, in particular L-lysine, L-
5 threonine, L-isoleucine and L-tryptophan, using coryneform bacteria in which at least the gnd gene is amplified.

Prior art

L-Amino acids are used in animal nutrition, in human medicine and in the pharmaceuticals industry.

10 It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular Corynebacterium glutamicum. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can
15 relate to fermentation measures, such as e.g. stirring and supply of oxygen, or the composition of the nutrient media, such as e.g. the sugar concentration during the fermentation, or the working up to the product form by e.g. ion exchange chromatography, or the intrinsic output
20 properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to
antimetabolites, such as e.g. the threonine analogue α -
25 amino- β -hydroxyvaleric acid (AHV), or are auxotrophic for metabolites of regulatory importance and produce L-amino acids such as e.g. threonine are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of
30 Corynebacterium glutamicum strains which produce L-amino acids.

Object of the invention

The inventors had the object of providing improved processes for the fermentative preparation of L-amino acids with coryneform bacteria.

5 Description of the invention

L-Amino acids are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and especially in animal nutrition. There is therefore a general interest in providing new improved processes for
10 the preparation of amino acids.

The invention provides a process for the fermentative preparation of L-amino acids, in particular L-lysine, L-threonine, L-isoleucine and L-tryptophan, using coryneform bacteria in which the nucleotide sequence which codes for
15 the enzyme 6-phosphogluconate dehydrogenase (EC number 1.1.1.44) (gnd gene) is amplified, in particular over-expressed.

The strains employed preferably already produce L-amino acids before amplification of the gnd gene.

20 Preferred embodiments are to be found in the claims.

The term "amplification" in this connection describes the increase in the intracellular activity of one or more enzymes in a microorganism which are coded by the corresponding DNA, for example by increasing the number of
25 copies of the gene or genes, using a potent promoter or using a gene which codes for a corresponding enzyme having a high activity, and optionally combining these measures.

The microorganisms which the present invention provides can prepare L-amino acids from glucose, sucrose, lactose,
30 fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They are representatives of

coryneform bacteria, in particular of the genus
Corynebacterium. Of the genus Corynebacterium, there may be
mentioned in particular the species Corynebacterium
glutamicum, which is known among experts for its ability to
5 produce L-amino acids.

Suitable strains of the genus Corynebacterium, in
particular of the species Corynebacterium glutamicum, are,
for example, the known wild-type strains

Corynebacterium glutamicum ATCC13032
10 Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
Corynebacterium thermoaminogenes FERM BP-1539
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869
15 Brevibacterium divaricatum ATCC14020

and L-amino acid-producing mutants prepared therefrom,

such as, for example, the L-threonine-producing strains

Corynebacterium glutamicum ATCC21649
Brevibacterium flavum BB69
20 Brevibacterium flavum DSM5399
Brevibacterium lactofermentum FERM-BP 269
Brevibacterium lactofermentum TBB-10

and such as, for example, the L-isoleucine-producing
strains

25 Corynebacterium glutamicum ATCC 14309
Corynebacterium glutamicum ATCC 14310
Corynebacterium glutamicum ATCC 14311
Corynebacterium glutamicum ATCC 15168
Corynebacterium ammoniagenes ATCC 6871

30 and such as, for example, the L-tryptophan-producing
strains

Corynebacterium glutamicum ATCC21850
Corynebacterium glutamicum KY9218 (pKW9901)

and such as, for example, the L-lysine-producing strains

Corynebacterium glutamicum FERM-P 1709

Brevibacterium flavum FERM-P 1708

Brevibacterium lactofermentum FERM-P 1712

5 *Corynebacterium glutamicum* FERM-P 6463

Corynebacterium glutamicum FERM-P 6464

Corynebacterium glutamicum DSM5715

Corynebacterium glutamicum DM58-1

Corynebacterium glutamicum DSM12866.

10 It has been found that coryneform bacteria produce L-amino acids, in particular L-lysine, L-threonine, L-isoleucine and L-tryptophan, in an improved manner after over-expression of the *gnd* gene which codes for 6-phosphogluconate dehydrogenase (EC number 1.1.1.44).

15 The *gnd* gene codes for the enzyme 6-phosphogluconate dehydrogenase, which catalyses the oxidative decarboxylation of 6-phosphogluconic acid to ribulose 5-phosphate. The nucleotide sequence of the *gnd* gene is disclosed in JP-A-9-224662. The *gnd* gene described in the
20 text reference mentioned is used according to the invention for the first time. Alleles of the *gnd* gene which result from the degeneracy of the genetic code or due to sense mutations of neutral function can furthermore be used.

To achieve an amplification (e.g. over-expression), the
25 number of copies of the corresponding genes is increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene is mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible
30 promoters, it is additionally possible to increase the expression in the course of fermentative L-amino acid formation. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of
35 the enzyme protein. The genes or gene constructs are either

present here in plasmids with a varying number of copies, or are integrated and amplified in the chromosome.

Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)),
10 in Eikmanns et al. (Gene 102, 93-98 (1991)), in European Patent Specification EPS 0 472 869, in US Patent 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of
15 Bacteriology 175, 1001-1007 (1993)), in Patent Application WO 96/15246, in Malumbres et al. (Gene 134, 15-24 (1993)), in Japanese Laid-Open Specification JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)) and in known textbooks of genetics and
20 molecular biology.

By way of example, 6-phosphogluconate dehydrogenase was over-expressed with the aid of a plasmid. The *E. coli* - *C. glutamicum* shuttle vector pEC-T18mob2 shown in Figure 1 was used for this. After incorporation of the *gnd* gene into the
25 *EcoRI* cleavage site of pEC-T18mob2, the plasmid pECgnd shown in Figure 2 was formed.

Other plasmid vectors which are capable of replication in *C. glutamicum*, such as e.g. pEKEx1 (Eikmanns et al., Gene 102:93-98 (1991)) or pZ8-1 (EP-B- 0 375 889), can be used
30 in the same way.

In addition, it may be advantageous for the production of L-amino acids to amplify one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the pentose phosphate pathway or of amino

acid export, in addition to amplification of the gnd gene which codes for 6-phosphogluconate dehydrogenase.

Thus, for example, in particular for the preparation of L-threonine, one or more genes chosen from the group

5 consisting of

- the hom gene which codes for homoserine dehydrogenase (Peoples et al., Molecular Microbiology 2, 63-72 (1988)) or the hom^{dr} allele which codes for a "feed back resistant" homoserine dehydrogenase (Archer et al., Gene
10 107, 53-59 (1991),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns et al., Journal of Bacteriology 174: 6076-6086 (1992)),
- the pyc gene which codes for pyruvate carboxylase
15 (Peters-Wendisch et al., Microbiology 144: 915-927 (1998)),
- the mqo gene which codes for malate:quinone oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
- 20 • the tkt gene which codes for transketolase (accession number AB023377 of the databank of European Molecular Biology Laboratories (EMBL, Heidelberg, Germany)),
- the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661),
- 25 • the thrE gene which codes for threonine export (DE 199 41 478.5; DSM 12840),
- the zwf gene (DE 199 59 328.0; DSM 13115),
- the eno gene which codes for enolase (DE: 199 41 478.5)

can be amplified, in particular over-expressed, at the same time.

Thus, for example, in particular for the preparation of L-lysine, one or more genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- a lysC gene which codes for a feed back resistant aspartate kinase (Kalinowski et al. (1990), Molecular and General Genetics 224: 317-324),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pyc gene which codes for pyruvate carboxylase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the mgo gene which codes for malate-quinone oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
- the tkt gene which codes for transketolase (accession number AB023377 of the databank of European Molecular Biologies Laboratories (EMBL, Heidelberg, Germany)),
- the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661),
- the lysE gene which codes for lysine export (DE-A-195 48 222),
- the zwf gene (DE 199 59 328.0; DSM 13115),
- the eno gene which codes for enolase (DE 199 47 791.4)

can be amplified, in particular over-expressed, at the same time.

It may furthermore be advantageous for the production of L-amino acids at the same time to attenuate one or more of
5 the genes chosen from the group consisting of:

- the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),
- 10 • the poxB gene which codes for pyruvate oxidase (DE 199 51 975.7; DSM 13114),
- the zwa2 gene (DE: 199 59 327.2; DSM 13113)

in addition to the amplification of the gnd gene.

In addition to over-expression of 6-phosphogluconate
15 dehydrogenase, it may furthermore be advantageous for the production of L-amino acids to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London,
20 UK, 1982).

The microorganisms prepared according to the invention can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for
25 the purpose of L-amino acid production. A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 30 1991)) or in the textbook by Storhas (Bioreaktoren und

periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular microorganisms in a suitable manner.

- 5 Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture. Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture. Potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

- Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of L-amino acid has formed. This target is usually reached within 10 hours to 160 hours.
- The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin derivatization, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190), or it can take place by reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The following microorganism has been deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

Escherichia coli K-12 DH5α/pEC-T18mob2 as DSM 13244

The following figures are attached:

- Figure 1: Map of the plasmid pEC-T18mob2
- Figure 2: Map of the plasmid pECgnd
- Figure 3: Map of the plasmid pBGNA
- 5 • Figure 4: Map of the plasmid pCR2.1poxBint

The base pair numbers stated are approx. values obtained in the context of reproducibility.

The abbreviations used have the following meaning:

Re Figure 1:

- | | | |
|----|-------------|---|
| 10 | Tet: | Resistance gene for tetracycline |
| | oriV: | Plasmid-coded replication origin of <i>E. coli</i> |
| | RP4mob: | mob region for mobilizing the plasmid |
| | rep: | Plasmid-coded replication origin from <i>C. glutamicum</i> plasmid pGA1 |
| 15 | per: | Gene for controlling the number of copies from pGA1 |
| | lacZ-alpha: | lacZ α gene fragment (N-terminus) of the β -Galactosidase gene |

Re Figure 2:

- | | | |
|----|------|---|
| 20 | Tet: | Resistance gene for tetracycline |
| | rep: | Plasmid-coded replication origin from <i>C. glutamicum</i> plasmid pGA1 |
| | per: | Gene for controlling the number of copies from pGA1 |
| 25 | lacZ | Cloning relict of the lacZ α gene fragment from pEC-T18mob2 |
| | gnd: | 6-Phosphogluconate dehydrogenase gene |

Re Figure 3:

- | | | |
|----|--------|---|
| | LacP: | Promoter of the <i>E. coli</i> lactose operon |
| 30 | CMV: | Promoter of cytomegalovirus |
| | ColE1: | Replication origin of the plasmid ColE1 |

TkpolyA: Polyadenylation site
Kan r: Kanamycin resistance gene
SV40ori: Replication origin of Simian virus 40
gnd: 6-Phosphogluconate dehydrogenase gene

5 Re Figure 4:

ColE1 ori: Replication origin of the plasmid ColE1
lacZ: Cloning relict of the lacZ α gene fragment
f1 ori: Replication origin of phage f1
KmR: Kanamycin resistance
10 ApR: Ampicillin resistance
poxBint: internal fragment of the poxB gene

Moreover, the following abbreviations have been used:

AccI: Cleavage site of the restriction enzyme AccI
BamHI: Cleavage site of the restriction enzyme BamHI
15 EcoRI: Cleavage site of the restriction enzyme EcoRI
HindIII: Cleavage site of the restriction enzyme HindIII
KpnI: Cleavage site of the restriction enzyme KpnI
PstI: Cleavage site of the restriction enzyme PstI
PvuI: Cleavage site of the restriction enzyme PvuI
20 SalI: Cleavage site of the restriction enzyme SalI
SacI: Cleavage site of the restriction enzyme SacI
SmaI: Cleavage site of the restriction enzyme SmaI
SphI: Cleavage site of the restriction enzyme SphI
XbaI: Cleavage site of the restriction enzyme XbaI
25 XhoI: Cleavage site of the restriction enzyme XhoI

Examples

The following examples will further illustrate this invention. The molecular biology techniques, e.g. plasmid DNA isolation, restriction enzyme treatment, ligations, standard transformations of *Escherichia coli* etc. used are, (unless stated otherwise), described by Sambrook et al., (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbour Laboratories, USA).

Example 1

- 10 Construction of a gene library of *Corynebacterium glutamicum* strain AS019

A DNA library of *Corynebacterium glutamicum* strain AS019 (Yoshihama et al., Journal of Bacteriology 162, 591-597 (1985)) was constructed using the Zap Express™ system, (Short et al., (1988) Nucleic Acids Research 16: 7583-7600), as described by O'Donohue (O'Donohue, M. (1997). The Cloning and Molecular Analysis of Four Common Aromatic Amino Acid Biosynthetic Genes from *Corynebacterium glutamicum*. Ph.D. Thesis, National University of Ireland, Galway). The Zap Express™ kit was purchased from Stratagene (Stratagene, 11011 North Torrey Pines Rd., La Jolla, California 92037) and used according to the manufacturers instructions. AS019-DNA was digested with restriction enzyme *Sau*3A and ligated to *Bam*HI treated and dephosphorylated Zap Express™ arms.

Example 2

Cloning and sequencing of the *gnd* gene

1. Construction of a *gnd* probe

A radiolabelled oligonucleotide, internal to the *gnd* gene, was used to probe the AS019 Zap Express™ library described above. The oligonucleotide was produced using degenerate PCR primers internal to the *gnd* gene. The

degenerate nucleotide primers designed for the PCR amplification of gnd DNA fragments were as follows:

gnd1: 5' ATG GTK CAC ACY GGY ATY GAR TA 3'

gnd2: 5' RGT CCA YTT RCC RGT RCC YTT 3'

5 with R=A+G; Y=C+T; K=T+G.

The estimated size of the resulting PCR product was 252 bp approximately.

Optimal PCR conditions were determined to be as follows:

35 cycles
10 94°C for 1 minute
55°C for 1 minute
72°C for 30 seconds
2.5 - 3.5 mM MgCl₂
100 - 150 ng AS019 genomic DNA

15 Sequence analysis of the resulting PCR product confirmed the product to be an internal portion of a gnd gene. Sequence analysis was carried out using the universal forward and reverse primers, and T7 sequencing kit from Pharmacia Biotech, (St. Albans, Herts, UK). The sequence of
20 the PCR product is shown in SEQ ID No. 1.

2. Cloning

Screening of the AS019 ÷ Zap Express™ library was carried out according to the ÷ Zap Express™ system protocol, (Stratagene, 11011 North Torrey Pines Rd., La Jolla,
25 California 92037). Southern Blot analysis was then carried out on isolated clones. Southern transfer of DNA was as described in the Schleicher and Schuell protocols manual employing Nytran™ as membrane („Nytran, Modified Nylon-66 Membrane Filters“ (March 1987), Schleicher and Schuell,
30 Dassel, Germany). Double stranded DNA fragments, generated using the same primers and optimal PCR conditions as described above, were radiolabelled with α -³²P-dCTP using

the MultiprimeTM DNA labelling kit from Amersham Life Science (Amersham Pharmacia Biotech UK Limited, Little Chalfont, Buckinghamshire, UK) according to the manufacturers instructions. Prehybridisation, hybridization and washing conditions were as described in the Schleicher and Schuell protocols manual. Autoradiography was carried out according to the procedure outlined in the handbook of Sambrook et al. using AgFa Curix RPIL film. Thus several gnd clones were identified. Plasmid DNA was isolated from one of the clones, designated pBGNA (Figure 3) and chosen for further analysis.

3. Sequencing

The Sanger Dideoxy chain termination method of Sanger et al. (Proceedings of the National Academy of Sciences USA 74, 5463-5467 (1977)) was used to sequence the cloned insert of pBGNA. The method was applied using the T7 sequencing kit and α -³⁵S-dCTP from Pharmacia Biotech (St. Albans, Herts, UK). Samples were electrophoresed for 3-8 hours on 6% polyacrylamide/urea gels in TBE buffer at a constant current of 50 mA, according to the Pharmacia cloning and sequencing instructions manual („T7 SequencingTM Kit“, ref.XY-010-00-19, Pharmacia Biotech, 1994). Sequence analysis was carried out using internal primers designed from the sequence known of the internal gnd PCR product (SEQ ID NO 1) allowing the entire gnd gene sequence to be deduced.

The sequences of the internal primers were as follows:

Internal primer 1: 5' GGT GGA TGC TGA AAC CG 3'
Internal primer 2: 5' GCT GCA TGC CTG CTG CG 3'
Internal primer 3: 5' TTG TTG CTT ACG CAC AG 3'
Internal primer 4: 5' TCG TAG GAC TTT GCT GG 3'

Sequence obtained was then analyzed using the DNA Strider programme, (Marck (1988), Nucleic Acids Research 16: 1829-1836), version 1.0 on an Apple Macintosh computer. This

program allowed for analyses such as restriction site usage, open reading frame analysis and codon usage determination. Searches between DNA sequence obtained and those in EMBL and Genbank databases were achieved using the
5 BLAST programme (Altschul et al., (1997), Nucleic Acids Research 25: 3389-3402). DNA and protein sequences were aligned using the Clustal V and Clustal W programs (Higgins and Sharp, 1988 Gene 73: 237-244).

The sequence thus obtained is shown in SEQ ID NO 2. The
10 analysis of the nucleotide sequence obtained revealed an open reading frame of 1377 base pairs which was designated as gnd gene. It codes for a protein of 459 amino acids shown in SEQ ID NO 3.

Example 3

15 Preparation of the shuttle vector pEC-T18mob2

The E. coli - C. glutamicum shuttle vector pEC-T18mob2 was constructed according to the prior art.

The vector contains the replication region rep of the plasmid pGA1 including the replication effector per (US-A-
20 5,175,108; Nesvera et al., Journal of Bacteriology 179, 1525-1532 (1997)), the tetracycline resistance-imparting tetA(2) gene of the plasmid pAG1 (US-A- 5,158,891; gene library entry at the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA) with accession number
25 AF121000), the replication region oriV of the plasmid pMB1 (Sutcliffe, Cold Spring Harbor Symposium on Quantitative Biology 43, 77-90 (1979)), the lacZ gene fragment including the lac promoter and a multiple cloning site (mcs) (Norlander et al. Gene 26, 101-106 (1983)) and the mob
30 region of the plasmid RP4 (Simon et al., (1983) Bio/Technology 1:784-791).

The vector constructed was transformed in the E. coli strain DH5 α (Hanahan, In: DNA cloning. A practical

approach. Vol. I. IRL-Press, Oxford, Washington DC, USA, 1985). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. 5 Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA, 1989), which had been supplemented with 5 mg/l tetracycline. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme 10 EcoRI and HindIII subsequent agarose gel electrophoresis (0.8%).

The plasmid was called pEC-T18mob2 and is shown in Figure 1. It is deposited in the form of the strain Escherichia coli K-12 strain DH5 α /pEC-T18mob2 at the Deutsche Sammlung 15 für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) as DSM 13244.

Example 4

Cloning of the gnd gene into the E. coli - C. glutamicum 20 shuttle vector pEC-T18mob2

PCR was used to amplify DNA fragments containing the entire gnd gene of C. glutamicum and flanking upstream and downstream regions using pBGNA as template. PCR reactions were carried out using oligonucleotide primers designed 25 from SEQ ID NO 2. The primers used were:

gnd fwd. primer: 5' ACT CTA GTC GGC CTA AAA TGG 3'
gnd rev. primer: 5' CAC ACA GGA AAC AGA TAT GAC 3'

PCR parameters were as follows:

35 cycles
95°C for 6 minutes
94°C for 1 minute
50°C for 1 minute
5 72°C for 45 seconds
1 mM MgCl₂
approx. 150-200ng pBGNA-DNA as template.

The PCR product obtained was cloned into the commercially available pGEM-T vector purchased from Promega Corp. (pGEM-
10 T Easy Vector System 1, cat. no. A1360, Promega UK, Southampton) using *E. coli* strain JM109 (Yanisch-Perron et al. *Gene*, 33: 103-119 (1985)) as a host. The entire *gnd* gene was subsequently isolated from the pGEM T-vector on an EcoRI fragment and cloned into the lacZ EcoRI site of the
15 *E. coli* - *C. glutamicum* shuttle vector pEC-T18mob2 (Figure 1), and designated pECgnd (Figure 2). Restriction enzyme analysis with AccI (Boehringer Mannheim GmbH, Germany) revealed the correct orientation (i. e. downstream the lac-Promotor) of the *gnd* gene in the lacZα gene of pEC-T18mob2.

20 Example 5

Preparation of amino acid producers with amplified 6-phosphogluconate dehydrogenase

Plasmid pECgnd from Example 3 was electroporated by the electroporation method of Tauch et al. (FEMS
25 Microbiological Letters, 123:343-347 (1994)) in the strains *Corynebacterium glutamicum* DSM 5399 and DSM 5714. The strain DSM 5399 is a threonine producer described in EP-B-0358940. The strain DSM 5714 is a lysine producer described in EP-B-0435132. Selection of transformants was carried out
30 by plating out the electroporation batch on LB agar (Sambrook et al., *Molecular cloning: a laboratory manual*. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with

25 mg/l kanamycin. The strains DSM5399/pECgnd and DSM5714/pECgnd were formed in this manner.

Example 6

Preparation of threonine

- 5 The *C. glutamicum* strain DSM5399/pECgnd obtained in Example 5 was cultured in a nutrient medium suitable for the production of threonine and the threonine content in the culture supernatant was determined.

10 For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). Brain-heart broth (Merck, Darmstadt, Germany) was used as the medium for the
15 preculture. Tetracycline (5 mg/l) was added to this medium. The preculture was incubated for 24 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1. The medium MM-threonine was used
20 for the main culture.

Medium MM-threonine:

CSL	5 g/l
MOPS	20 g/l
Glucose (autoclaved separately)	50g/l

Salts:

$(\text{NH}_4)_2\text{SO}_4$	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
CaCO_3	25 g/l

The CSL (corn steep liquor), MOPS
(morpholinopropanesulfonic acid) and the salt solution were
5 brought to pH 7 with aqueous ammonia and autoclaved. The
sterile substrate and vitamin solutions were then added, as
well as the CaCO_3 autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in a 100 ml
conical flask with baffles. Tetracycline (5 mg/l) was
10 added. Culturing was carried out at 33°C and 80%
atmospheric humidity.

After 48 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The concentration of threonine formed was determined with an amino acid analyzer from
5 Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD(660)	L-Threonin g/l
DSM5399/pECgnd	11.9	1.29
DSM5399	11.8	0.33

10

Example 7

Preparation of lysine

The C. glutamicum strain DSM5714/pECgnd obtained in Example
5 was cultured in a nutrient medium suitable for the
15 production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from
20 this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl 2.5 g/l

Bacto-Peptone 10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) was added to this medium. The preculture was incubated for 24 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main
5 culture was 0.05. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
$(\text{NH}_4)_2\text{SO}_4$	
KH_2PO_4	25 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	0.1 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	1.0 g/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	10 mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO_3	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO_3 autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 48 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, München). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-
5 BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 2.

Table 2

Strain	OD(660)	Lysine HCl g/l
DSM5715/pECgnd	7.7	14.7
DSM5715	7.1	13.7

10 Example 8

Preparation of a genomic cosmid gene library from
Corynebacterium glutamicum ATCC 13032.

Chromosomal DNA from Corynebacterium glutamicum ATCC 13032 was isolated as described by Tauch et al., (1995, Plasmid
15 33:168-179), and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product
20 Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vektor Kit, Code no. 251301) was cleaved
25 with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-

0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase. The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04).

5 The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of

10 Gigapack II XL Packing Extracts (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217). For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an

15 aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology 1:190) + 100 µg/ml

20 ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 9

Isolation and sequencing of the poxB gene

The cosmid DNA of an individual colony (Example 8) was

25 isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-

30 02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were

35 isolated with the QiaExII Gel Extraction Kit (Product No.

20021, Qiagen, Hilden, Germany). The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, Holland, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 μ g/ml zeocin. The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain-stopping method of Sanger et al. (1977, Proceedings of the National Academies of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. Al24.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZero1 derivatives were assembled to a continuous contig. The computer-assisted coding region analysis were prepared

with the XNIP program (Staden, 1986, Nucleic Acids Research 14:217-231). Further analyses were carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research 25:3389-3402), against the non-redundant
5 databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The resulting nucleotide sequence is shown in SEQ ID No. 4. Analysis of the nucleotide sequence showed an open reading frame of 1737 base pairs, which was called the poxB gene.
10 The poxB gene codes for a polypeptide of 579 amino acids (SEQ ID NO. 5).

Example 10

Preparation of an integration vector for integration mutagenesis of the poxB gene

15 From the strain ATCC 13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)). On the basis of the sequence of the poxB gene known for C. glutamicum from Example 9, the following oligonucleotides were chosen for the polymerase chain
20 reaction:

poxBint1:

5' TGC GAG ATG GTG AAT GGT GG 3'

poxBint2:

5' GCA TGA GGC AAC GCA TTA GC 3'

25 The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR protocols. A guide to methods and applications, 1990, Academic Press) with Pwo-Polymerase from Boehringer. With the aid of the
30 polymerase chain reaction, a DNA fragment approx. 0.9 kb in size was isolated, this carrying an internal fragment of the poxB gene and being shown in SEQ ID No. 6.

The amplified DNA fragment was ligated with the TOPO TA Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K4500-01) in the vector pCR2.1-TOPO (Mead et al. (1991) Bio/Technology 9:657-663). The E. coli Stamm DH5 α was then electroporated with the ligation batch (Hanahan, In: DNA cloning. A practical approach. Vol.I. IRL-Press, Oxford, Washington DC, USA, 1985). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 25 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1poxBint (Figure 4).

Plasmid pCR2.1poxBint has been deposited in the form of the strain Escherichia coli DH5 α /pCR2.1poxBint as DSM 13114 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty.

Example 11

Integration mutagenesis of the poxB gene in the lysine producer DSM 5715

The vector pCR2.1poxBint mentioned in Example 10 was electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) in Corynebacterium glutamicum DSM 5715. Strain DSM 5715 is an AEC-resistant lysine producer. The vector pCR2.1poxBint cannot replicate independently in DSM5715 and is retained in the cell only if it has integrated into the chromosome of DSM 5715. Selection of clones with pCR2.1poxBint

integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which
5 had been supplemented with 15 mg/l kanamycin. For detection of the integration, the poxBint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993).
10 Chromosomal DNA of a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes SalI, SacI and HindIII. The fragments formed were separated by agarose gel electrophoresis and hybridized at
15 68°C with the Dig hybridization kit from Boehringer. The plasmid pCR2.1poxBint mentioned in Example 9 had been inserted into the chromosome of DSM5715 within the chromosomal poxB gene. The strain was called DSM5715::pCR2.1poxBint.

20 Example 12

Effect of over-expression of the gnd gene with simultaneous elimination of the poxB gene on the preparation of lysine

12.1 Preparation of the strain

DSM5715::pCR2.1poxBint/pECgnd

25 The strain DSM5715::pCR2.1poxBint was transformed with the plasmid pECgnd using the electroporation method described by Liebl et al., (FEMS Microbiology Letters, 53:299-303 (1989)). Selection of the transformants took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M
30 sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-agar, which had been supplemented with 5 mg/l tetracycline and 25 mg/l kanamycin. Incubation was carried out for 2 days at 33°C.

Plasmid DNA was isolated in each case from a transformant by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915 -927), cleaved with the restriction endonuclease AccI, and the plasmid was checked by
5 subsequent agarose gel electrophoresis. The strain obtained in this way was called DSM5715:pCR2.1poxBint/pECgnd.

12.2 Preparation of L-lysine

The *C. glutamicum* strain DSM5715::pCR2.1poxBint/pECgnd
10 obtained in Example 12.1 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with
15 tetracycline (5 mg/l) and kanamycin (25 mg/l)) for 24 hours at 33°C. The cultures of the comparison strains were supplemented according to their resistance to antibiotics. Starting from this agar plate culture, a preculture was
seeded (10 ml medium in a 100 ml conical flask). The
20 complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
------	---------

Bacto-Peptone	10 g/l
---------------	--------

Bacto-Yeast extract	10 g/l
---------------------	--------

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) and kanamycin (25 mg/l) were added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded

from this preculture such that the initial OD (660nm) of the main culture was 0.1. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	58 g/l
$(\text{NH}_4)_2\text{SO}_4$	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO_3	25 g/l

5

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO_3 autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) and kanamycin (25 mg/l) were added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- 5 After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, München). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography
10 and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 3.

Table 3

Strain	OD	L-Lysine HCl g/l
DSM5715	10.8	16.0
DSM5715/pECgnd	7.6	16.5
DSM5715::pCR2.1poxBint	7.1	16.7
DSM5715::pCR2.1poxBint/ pECgnd	7.2	17.1

Patent claims

1. A process for the preparation of L-amino acids by
fermentation of coryneform bacteria
5 w h i c h c o m p r i s e s
 carrying out the following steps:
 - a) fermentation of the desired L-amino acid-
 producing bacteria in which at least the gnd gene
 is amplified,
 - 10 b) concentration of the L-amino acid in the medium
 or in the cells of the bacteria and
 - c) isolation of the L-amino acid produced.
2. The process as claimed in claim 1,
 w h e r e i n
15 bacteria in which further genes of the biosynthesis
 pathway of the desired L-amino acid are additionally
 amplified, in particular over-expressed, are employed.
3. The process as claimed in claim 1,
 w h e r e i n
20 coryneform bacteria which prepare L-threonine, L-
 lysine, L-isoleucine or L-tryptophan are used.
4. The process as claimed in claim 3,
 w h e r e i n
 coryneform bacteria which prepare L-lysine are used.
- 25 5. A process for the fermentative preparation of L-lysine
 as claimed in claim 2,
 w h e r e i n
 in the coryneform microorganisms which in particular
 already produce L-lysine, one or more genes chosen from
30 the group consisting of
 - 5.1 the dapA gene which codes for
 dihydrodipicolinate synthase,

- 5.2 the lysC gene which codes for a feed back resistant aspartate kinase,
- 5.3 the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase,
- 5 5.4 the pyc gene which codes for pyruvate carboxylase,
- 5.5 the tkt gene which codes for transketolase,
- 5.6 the zwf gene which codes for glucose 6-phosphate dehydrogenase,
- 10 5.7 the lysE gene which codes for lysine export,
- 5.8 the zwal gene
- 5.9 the eno gene which codes for enolase
- is or are amplified, in particular over-expressed, at the same time.
- 15 6. A process for the fermentative preparation of L-threonine as claimed in claim 2, where in
- in the coryneform microorganisms which in particular already produce L-threonine, one or more genes chosen
- 20 from the group consisting of
- 6.1 the hom gene which codes for homoserine dehydrogenase or the hom^{dr} allele which codes for a "feed back resistant" homoserine dehydrogenase,
- 6.2 the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase,
- 25 6.3 the pyc gene which codes for pyruvate carboxylase,

6.4 the mqo gene which codes for malate:quinone
oxidoreductase,

6.5 the tkt gene which codes for transketolase,

6.6 the zwf gene which codes for glucose 6-phosphate
dehydrogenase,

6.7 the thrE gene which codes for threonine export,

6.8 the zwal gene,

6.9 the eno gene which codes for enolase

is or are amplified, in particular over-expressed, at
the same time.

7. The process as claimed in claim 2,

wherein

for the preparation of L-amino acids, in particular L-
lysine or L-threonine, bacteria in which one or more
genes chosen from the group consisting of,

7.1 the pck gene which codes for phosphoenol pyruvate
carboxykinase

7.2 the pgi gene which codes for glucose 6-phosphate
isomerase

7.3 the poxB gene which codes for pyruvate oxidase or

7.4 the zwa2 gene

is or are attenuated at the same time, are fermented.

8. The process as claimed in claims 2 to 6,

wherein

to achieve the amplification, the number of copies of
the genes or nucleotide sequences is increased by
transformation of the microorganisms with plasmid

vectors which carry these genes or nucleotide sequences.

9. The plasmid vector pEC-T18mob2 deposited under the designation DSM 13244 in *E. coli* K-12 DH5 α , shown in
5 Figure 1.
10. A coryneform microorganism, in particular of the genus *Corynebacterium*, transformed by the introduction of the plasmid vector as claimed in claim 9, which additionally contains the *gnd* gene.

Figure 1: Map of the plasmid pEC-T18mob2




Figure 2: Map of the plasmid pECgnd




Figure 3: Map of the plasmid pBGNA





Figure 4: Map of the plasmid pCR2.1poxBint



SEQUENCE PROTOCOL

<110> National University of Ireland, Galway
 Degussa-Hüls AG

5 <120> Process for the fermentative preparation of
 L-amino acids using coryneform bacteria.

10 <130> 990229 BT

<140>
 <141>

15 <160> 6

<170> PatentIn Ver. 2.1

<210> 1
 <211> 252
 20 <212> DNA
 <213> Corynebacterium glutamicum

<400> 1
 atggtccaca acggcatcga gtacgccgac atgcagggtca tcggcgaggc ataccacctt 60
 25 ctgccctacg cagcaggcat gcagccagct gaaatcgctg aggttttcaa ggaatggaac 120
 gcaggcgacc tggattccta cctcatcgaa atcaccgcag aggttctctc ccagggtggat 180
 gctgaaaccg gcaagccact aatcgacgtc atcgttgacg ctgcagggtca gaagggcacc 240
 ggcaagtgga ct 252

30 <210> 2
 <211> 2335
 <212> DNA
 <213> Corynebacterium glutamicum

35 <220>
 <221> CDS
 <222> (474)..(1850)
 <223> gnd

40 <400> 2
 ttgttcggcc acgatgacac cggagctcac agcagaaatg aagtcggtgt tgttgttgat 60
 gccgacgacc atttttccag gggcggaat catgctggcg actgatccag tggattcggc 120
 45 gatggcggcg tagacaccac cgttgaccaa gccaccact tgcagggtgt tggatgccac 180
 gtgaagtctg ctgaccaccc ggccgggctc gatggtggtg tagcgcagcc ccagattgcy 240
 50 gtcgaggcca taattggcgt tgttgagtgc ttcaagttcg tctgtggtta aagctctggt 300
 ggcggcaagt tctgcaagcg aaagcagatc ttggggttga tcatcgcggg aagtcataat 360
 taattactct agtcggccta aaatggttgg attttcacct cctgtgacct ggtaaaatcg 420
 55 ccactacccc caaatggtca caccttttag gccgattttg ctgacaccgg gct atg 476
 Met
 1

60 ccg tca agt acg atc aat aac atg act aat gga gat aat ctc gca cag 524
 Pro Ser Ser Thr Ile Asn Asn Met Thr Asn Gly Asp Asn Leu Ala Gln

5 10 15

	atc ggc gtt gta ggc cta gca gta atg ggc tca aac ctc gcc cgc aac	572
	Ile Gly Val Val Gly Leu Ala Val Met Gly Ser Asn Leu Ala Arg Asn	
	20 25 30	
5	ttc gcc cgc aac ggc aac act gtc gct gtc tac aac cgc agc act gac	620
	Phe Ala Arg Asn Gly Asn Thr Val Ala Val Tyr Asn Arg Ser Thr Asp	
	35 40 45	
10	aaa acc gac aag ctc atc gcc gat cac ggc tcc gaa ggc aac ttc atc	668
	Lys Thr Asp Lys Leu Ile Ala Asp His Gly Ser Glu Gly Asn Phe Ile	
	50 55 60 65	
15	cct tct gca acc gtc gaa gag ttc gta gca tcc ctg gaa aag cca cgc	716
	Pro Ser Ala Thr Val Glu Glu Phe Val Ala Ser Leu Glu Lys Pro Arg	
	70 75 80	
20	cgc gcc atc atc atg gtt cag gct ggt aac gcc acc gac gca gtc atc	764
	Arg Ala Ile Ile Met Val Gln Ala Gly Asn Ala Thr Asp Ala Val Ile	
	85 90 95	
25	aac cag ctg gca gat gcc atg gac gaa ggc gac atc atc atc gac ggc	812
	Asn Gln Leu Ala Asp Ala Met Asp Glu Gly Asp Ile Ile Ile Asp Gly	
	100 105 110	
30	ggc aac gcc ctc tac acc gac acc att cgt cgc gag aag gaa atc tcc	860
	Gly Asn Ala Leu Tyr Thr Asp Thr Ile Arg Arg Glu Lys Glu Ile Ser	
	115 120 125	
35	gca cgc ggt ctc cac ttc gtc ggt gct ggt atc tcc ggc ggc gaa gaa	908
	Ala Arg Gly Leu His Phe Val Gly Ala Gly Ile Ser Gly Gly Glu Glu	
	130 135 140 145	
40	ggc gca ctc aac ggc cca tcc atc atg cct ggt ggc cca gca aag tcc	956
	Gly Ala Leu Asn Gly Pro Ser Ile Met Pro Gly Gly Pro Ala Lys Ser	
	150 155 160	
45	tac gag tcc ctc gga cca ctg ctt gag tcc atc gct gcc aac gtt gac	1004
	Tyr Glu Ser Leu Gly Pro Leu Leu Glu Ser Ile Ala Ala Asn Val Asp	
	165 170 175	
50	ggc acc cca tgt gtc acc cac atc ggc cca gac ggc gcc ggc cac ttc	1052
	Gly Thr Pro Cys Val Thr His Ile Gly Pro Asp Gly Ala Gly His Phe	
	180 185 190	
55	gtc aag atg gtc cac aac ggc atc gag tac gcc gac atg cag gtc atc	1100
	Val Lys Met Val His Asn Gly Ile Glu Tyr Ala Asp Met Gln Val Ile	
	195 200 205	
60	ggc gag gca tac cac ctt ctg ccc tac gca gca ggc atg cag cca gct	1148
	Gly Glu Ala Tyr His Leu Leu Pro Tyr Ala Ala Gly Met Gln Pro Ala	
	210 215 220 225	
65	gaa atc gct gag gtt ttc aag gaa tgg aac gca ggc gac ctg gat tcc	1196
	Glu Ile Ala Glu Val Phe Lys Glu Trp Asn Ala Gly Asp Leu Asp Ser	
	230 235 240	
70	tac ctc atc gaa atc acc gca gag gtt ctc tcc cag gtg gat gct gaa	1244
	Tyr Leu Ile Glu Ile Thr Ala Glu Val Leu Ser Gln Val Asp Ala Glu	
	245 250 255	
75	acc ggc aag cca cta atc gac gtc atc gtt gac gct gca ggt cag aag	1292
	Thr Gly Lys Pro Leu Ile Asp Val Ile Val Asp Ala Ala Gly Gln Lys	
	260 265 270	

5	ggc acc ggc aag tgg act gtc aag gct gct ctt gat ctg ggt att gct 1340 Gly Thr Gly Lys Trp Thr Val Lys Ala Ala Leu Asp Leu Gly Ile Ala 275 280 285
	acc acc ggc atc ggc gaa cgt gtt ttc gca cgt gca ctc tcc ggc gca 1388 Thr Thr Gly Ile Gly Glu Arg Val Phe Ala Arg Ala Leu Ser Gly Ala 290 295 300 305
10	acc agc cag cgc gct gca gca cag ggc aac cta cct gca ggt gtc ctc 1436 Thr Ser Gln Arg Ala Ala Ala Gln Gly Asn Leu Pro Ala Gly Val Leu 310 315 320
15	acc gat ctg gaa gca ctt ggc gtg gac aag gca cag ttc gtc gaa gga 1484 Thr Asp Leu Glu Ala Leu Gly Val Asp Lys Ala Gln Phe Val Glu Gly 325 330 335
20	ctt cgc cgt gca ctg tac gca tcc aag ctt gtt gct tac gca cag ggc 1532 Leu Arg Arg Ala Leu Tyr Ala Ser Lys Leu Val Ala Tyr Ala Gln Gly 340 345 350
25	ttc gac gag atc aag gct ggc tcc gac gag aac aac tgg gac gtt gac 1580 Phe Asp Glu Ile Lys Ala Gly Ser Asp Glu Asn Asn Trp Asp Val Asp 355 360 365
30	cct cgc gac ctc gct acc atc tgg cgc ggc ggc tgc atc att cgc gct 1628 Pro Arg Asp Leu Ala Thr Ile Trp Arg Gly Gly Cys Ile Ile Arg Ala 370 375 380 385
35	aag ttc ctc aac cgc atc gtc gaa gca tac gat gca aac gct gaa ctt 1676 Lys Phe Leu Asn Arg Ile Val Glu Ala Tyr Asp Ala Asn Ala Glu Leu 390 395 400
40	gag tcc ctg ctg ctc gat cct tac ttc aag agc gag ctc ggc gac ctc 1724 Glu Ser Leu Leu Leu Asp Pro Tyr Phe Lys Ser Glu Leu Gly Asp Leu 405 410 415
45	atc gat tca tgg cgt cgc gtg att gtc acc gcc acc cag ctt ggc ctg 1772 Ile Asp Ser Trp Arg Arg Val Ile Val Thr Ala Thr Gln Leu Gly Leu 420 425 430
50	cca atc cca gtg ttc gct tcc tcc ctg tcc tac tac gac agc ctg cgt 1820 Pro Ile Pro Val Phe Ala Ser Ser Leu Ser Tyr Tyr Asp Ser Leu Arg 435 440 445
55	gca gag cgt ctg cca gca gcc ctg atc cac tagtgtcgac ctgcaggcgc 1870 Ala Glu Arg Leu Pro Ala Ala Leu Ile His 450 455
60	gcgagctcca gcttttggtc ccttttagtga gggttaattt cgagcttggc gtaatcaagg 1930 tcatagtctgt ttctgtgtg aaattgttat ccgctcaciaa ttccacaciaa tatacgagcc 1990 ggaagtataa agtgtaaagc ctggggtgcc taatgagtga gctaactcac agtaattgcg 2050 gctagcggat ctgacgggtc actaaaccag ctctgcttat atagacctcc caccgtacac 2110 gcctaccgcc catttgcgtc aatggggcgg agttgttacg acattttgga aagtcccgtt 2170 gattttggtg ccaaaacaaa ctcccattga cgtcaatggg gtggagactt ggaaatcccc 2230 gtgagtcaaa ccgctatcca cgcccattga tgtactgcca aaaccgcac accatggtaa 2290

tagcgatgac taatacgtag atgtactgcc aagtaggaaa gtccc

2335

5 <210> 3
 <211> 459
 <212> PRT
 <213> Corynebacterium glutamicum

 <400> 3
 10 Met Pro Ser Ser Thr Ile Asn Asn Met Thr Asn Gly Asp Asn Leu Ala
 1 5 10 15
 Gln Ile Gly Val Val Gly Leu Ala Val Met Gly Ser Asn Leu Ala Arg
 20 25 30
 15 Asn Phe Ala Arg Asn Gly Asn Thr Val Ala Val Tyr Asn Arg Ser Thr
 35 40 45
 20 Asp Lys Thr Asp Lys Leu Ile Ala Asp His Gly Ser Glu Gly Asn Phe
 50 55 60
 Ile Pro Ser Ala Thr Val Glu Glu Phe Val Ala Ser Leu Glu Lys Pro
 65 70 75 80
 25 Arg Arg Ala Ile Ile Met Val Gln Ala Gly Asn Ala Thr Asp Ala Val
 85 90 95
 Ile Asn Gln Leu Ala Asp Ala Met Asp Glu Gly Asp Ile Ile Ile Asp
 100 105 110
 30 Gly Gly Asn Ala Leu Tyr Thr Asp Thr Ile Arg Arg Glu Lys Glu Ile
 115 120 125
 35 Ser Ala Arg Gly Leu His Phe Val Gly Ala Gly Ile Ser Gly Gly Glu
 130 135 140
 Glu Gly Ala Leu Asn Gly Pro Ser Ile Met Pro Gly Gly Pro Ala Lys
 145 150 155 160
 40 Ser Tyr Glu Ser Leu Gly Pro Leu Leu Glu Ser Ile Ala Ala Asn Val
 165 170 175
 Asp Gly Thr Pro Cys Val Thr His Ile Gly Pro Asp Gly Ala Gly His
 180 185 190
 45 Phe Val Lys Met Val His Asn Gly Ile Glu Tyr Ala Asp Met Gln Val
 195 200 205
 50 Ile Gly Glu Ala Tyr His Leu Leu Pro Tyr Ala Ala Gly Met Gln Pro
 210 215 220
 Ala Glu Ile Ala Glu Val Phe Lys Glu Trp Asn Ala Gly Asp Leu Asp
 225 230 235 240
 55 Ser Tyr Leu Ile Glu Ile Thr Ala Glu Val Leu Ser Gln Val Asp Ala
 245 250 255
 Glu Thr Gly Lys Pro Leu Ile Asp Val Ile Val Asp Ala Ala Gly Gln
 260 265 270
 60 Lys Gly Thr Gly Lys Trp Thr Val Lys Ala Ala Leu Asp Leu Gly Ile
 275 280 285

Ala Thr Thr Gly Ile Gly Glu Arg Val Phe Ala Arg Ala Leu Ser Gly
 290 295 300
 5 Ala Thr Ser Gln Arg Ala Ala Ala Gln Gly Asn Leu Pro Ala Gly Val
 305 310 315 320
 Leu Thr Asp Leu Glu Ala Leu Gly Val Asp Lys Ala Gln Phe Val Glu
 325 330 335
 10 Gly Leu Arg Arg Ala Leu Tyr Ala Ser Lys Leu Val Ala Tyr Ala Gln
 340 345 350
 Gly Phe Asp Glu Ile Lys Ala Gly Ser Asp Glu Asn Asn Trp Asp Val
 355 360 365
 15 Asp Pro Arg Asp Leu Ala Thr Ile Trp Arg Gly Gly Cys Ile Ile Arg
 370 375 380
 20 Ala Lys Phe Leu Asn Arg Ile Val Glu Ala Tyr Asp Ala Asn Ala Glu
 385 390 395 400
 Leu Glu Ser Leu Leu Leu Asp Pro Tyr Phe Lys Ser Glu Leu Gly Asp
 405 410 415
 25 Leu Ile Asp Ser Trp Arg Arg Val Ile Val Thr Ala Thr Gln Leu Gly
 420 425 430
 Leu Pro Ile Pro Val Phe Ala Ser Ser Leu Ser Tyr Tyr Asp Ser Leu
 435 440 445
 30 Arg Ala Glu Arg Leu Pro Ala Ala Leu Ile His
 450 455
 35 <210> 4
 <211> 2160
 <212> DNA
 <213> Corynebacterium glutamicum
 40 <220>
 <221> CDS
 <222> (327)..(2063)
 <223> poxB
 45 <400> 4
 ttagaggcga ttctgtgagg tcactttttg tggggtcggg gtctaaattt ggccagtttt 60
 cgaggcgacc agacaggcgt gcccacgatg tttaaataagg cgatcggtgg gcatctgtgt 120
 50 ttggtttcga cgggctgaaa ccaaaccaga ctgccagca acgacggaaa tcccaaaagt 180
 gggcatccct gtttggtacc gagtaccac cggggcctga aactccctgg caggcgggcg 240
 55 aagcgtggca acaactggaa tttaagagca caattgaagt cgcaccaagt taggcaacac 300
 aatagccata acgttgagga gttcag atg gca cac agc tac gca gaa caa tta 353
 Met Ala His Ser Tyr Ala Glu Gln Leu
 1 5
 60 att gac act ttg gaa gct caa ggt gtg aag cga att tat ggt ttg gtg 401
 Ile Asp Thr Leu Glu Ala Gln Gly Val Lys Arg Ile Tyr Gly Leu Val
 10 15 20 25

	ggt gac agc ctt aat ccg atc gtg gat gct gtc cgc caa tca gat att	449
	Gly Asp Ser Leu Asn Pro Ile Val Asp Ala Val Arg Gln Ser Asp Ile	
	30 35 40	
5	gag tgg gtg cac gtt cga aat gag gaa gcg gcg gcg ttt gca gcc ggt	497
	Glu Trp Val His Val Arg Asn Glu Glu Ala Ala Ala Phe Ala Ala Gly	
	45 50 55	
10	gcg gaa tcg ttg atc act ggg gag ctg gca gta tgt gct gct tct tgt	545
	Ala Glu Ser Leu Ile Thr Gly Glu Leu Ala Val Cys Ala Ala Ser Cys	
	60 65 70	
15	ggt cct gga aac aca cac ctg att cag ggt ctt tat gat tcg cat cga	593
	Gly Pro Gly Asn Thr His Leu Ile Gln Gly Leu Tyr Asp Ser His Arg	
	75 80 85	
20	aat ggt gcg aag gtg ttg gcc atc gct agc cat att ccg agt gcc cag	641
	Asn Gly Ala Lys Val Leu Ala Ile Ala Ser His Ile Pro Ser Ala Gln	
	90 95 100 105	
25	att ggt tcg acg ttc ttc cag gaa acg cat ccg gag att ttg ttt aag	689
	Ile Gly Ser Thr Phe Phe Gln Glu Thr His Pro Glu Ile Leu Phe Lys	
	110 115 120	
30	gaa tgc tct ggt tac tgc gag atg gtg aat ggt ggt gag cag ggt gaa	737
	Glu Cys Ser Gly Tyr Cys Glu Met Val Asn Gly Gly Glu Gln Gly Glu	
	125 130 135	
35	cgc att ttg cat cac gcg att cag tcc acc atg gcg ggt aaa ggt gtg	785
	Arg Ile Leu His His Ala Ile Gln Ser Thr Met Ala Gly Lys Gly Val	
	140 145 150	
40	tcg gtg gta gtg att cct ggt gat atc gct aag gaa gac gca ggt gac	833
	Ser Val Val Val Ile Pro Gly Asp Ile Ala Lys Glu Asp Ala Gly Asp	
	155 160 165	
45	ggt act tat tcc aat tcc act att tct tct ggc act cct gtg gtg ttc	881
	Gly Thr Tyr Ser Asn Ser Thr Ile Ser Ser Gly Thr Pro Val Val Phe	
	170 175 180 185	
50	ccg gat cct act gag gct gca gcg ctg gtg gag gcg att aac aac gct	929
	Pro Asp Pro Thr Glu Ala Ala Ala Leu Val Glu Ala Ile Asn Asn Ala	
	190 195 200	
55	aag tct gtc act ttg ttc tgc ggt gcg ggc gtg aag aat gct cgc gcg	977
	Lys Ser Val Thr Leu Phe Cys Gly Ala Gly Val Lys Asn Ala Arg Ala	
	205 210 215	
60	cag gtg ttg gag ttg gcg gag aag att aaa tca ccg atc ggg cat gcg	1025
	Gln Val Leu Glu Leu Ala Glu Lys Ile Lys Ser Pro Ile Gly His Ala	
	220 225 230	
65	ctg ggt ggt aag cag tac atc cag cat gag aat ccg ttt gag gtc ggc	1073
	Leu Gly Gly Lys Gln Tyr Ile Gln His Glu Asn Pro Phe Glu Val Gly	
	235 240 245	
70	atg tct ggc ctg ctt ggt tac ggc gcc tgc gtg gat gcg tcc aat gag	1121
	Met Ser Gly Leu Leu Gly Tyr Gly Ala Cys Val Asp Ala Ser Asn Glu	
	250 255 260 265	
75	gcg gat ctg ctg att cta ttg ggt acg gat ttc cct tat tct gat ttc	1169
	Ala Asp Leu Leu Ile Leu Leu Gly Thr Asp Phe Pro Tyr Ser Asp Phe	

	270								275				280				
5	ctt	cct	aaa	gac	aac	gtt	gcc	cag	gtg	gat	atc	aac	ggt	gcg	cac	att	1217
	Leu	Pro	Lys	Asp	Asn	Val	Ala	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	
				285					290					295			
10	ggt	cga	cgt	acc	acg	gtg	aag	tat	ccg	gtg	acc	ggt	gat	gtt	gct	gca	1265
	Gly	Arg	Arg	Thr	Thr	Val	Lys	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	
			300					305					310				
15	aca	atc	gaa	aat	att	ttg	cct	cat	gtg	aag	gaa	aaa	aca	gat	cgt	tcc	1313
	Thr	Ile	Glu	Asn	Ile	Leu	Pro	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	
		315					320					325					
20	ttc	ctt	gat	cgg	atg	ctc	aag	gca	cac	gag	cgt	aag	ttg	agc	tcg	gtg	1361
	Phe	Leu	Asp	Arg	Met	Leu	Lys	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	
	330					335					340					345	
25	gta	gag	acg	tac	aca	cat	aac	gtc	gag	aag	cat	gtg	cct	att	cac	cct	1409
	Val	Glu	Thr	Tyr	Thr	His	Asn	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	
					350					355					360		
30	gaa	tac	gtt	gcc	tct	att	ttg	aac	gag	ctg	gcg	gat	aag	gat	gcg	gtg	1457
	Glu	Tyr	Val	Ala	Ser	Ile	Leu	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	
				365					370					375			
35	ttt	act	gtg	gat	acc	ggc	atg	tgc	aat	gtg	tgg	cat	gcg	agg	tac	atc	1505
	Phe	Thr	Val	Asp	Thr	Gly	Met	Cys	Asn	Val	Trp	His	Ala	Arg	Tyr	Ile	
			380					385					390				
40	gag	aat	ccg	gag	gga	acg	cgc	gac	ttt	gtg	ggt	tca	ttc	cgc	cac	ggc	1553
	Glu	Asn	Pro	Glu	Gly	Thr	Arg	Asp	Phe	Val	Gly	Ser	Phe	Arg	His	Gly	
		395					400					405					
45	acg	atg	gct	aat	gcg	ttg	cct	cat	gcg	att	ggt	gcg	caa	agt	gtt	gat	1601
	Thr	Met	Ala	Asn	Ala	Leu	Pro	His	Ala	Ile	Gly	Ala	Gln	Ser	Val	Asp	
	410					415					420					425	
50	cga	aac	cgc	cag	gtg	atc	gcg	atg	tgt	ggc	gat	ggt	ggt	ttg	ggc	atg	1649
	Arg	Asn	Arg	Gln	Val	Ile	Ala	Met	Cys	Gly	Asp	Gly	Gly	Leu	Gly	Met	
					430					435					440		
55	ctg	ctg	ggt	gag	ctt	ctg	acc	gtt	aag	ctg	cac	caa	ctt	ccg	ctg	aag	1697
	Leu	Leu	Gly	Glu	Leu	Leu	Thr	Val	Lys	Leu	His	Gln	Leu	Pro	Leu	Lys	
				445					450					455			
60	gct	gtg	gtg	ttt	aac	aac	agt	tct	ttg	ggc	atg	gtg	aag	ttg	gag	atg	1745
	Ala	Val	Val	Phe	Asn	Asn	Ser	Ser	Leu	Gly	Met	Val	Lys	Leu	Glu	Met	
			460					465					470				
65	ctc	gtg	gag	gga	cag	cca	gaa	ttt	ggt	act	gac	cat	gag	gaa	gtg	aat	1793
	Leu	Val	Glu	Gly	Gln	Pro	Glu	Phe	Gly	Thr	Asp	His	Glu	Glu	Val	Asn	
			475				480					485					
70	ttc	gca	gag	att	gcg	gcg	gct	gcg	ggt	atc	aaa	tcg	gta	cgc	atc	acc	1841
	Phe	Ala	Glu	Ile	Ala	Ala	Ala	Ala	Gly	Ile	Lys	Ser	Val	Arg	Ile	Thr	
	490					495					500					505	
75	gat	ccg	aag	aaa	gtt	cgc	gag	cag	cta	gct	gag	gca	ttg	gca	tat	cct	1889
	Asp	Pro	Lys	Lys	Val	Arg	Glu	Gln	Leu	Ala	Glu	Ala	Leu	Ala	Tyr	Pro	
					510					515					520		
80	gga	cct	gta	ctg	atc	gat	atc	gtc	acg	gat	cct	aat	gcg	ctg	tcg	atc	1937

Gly Pro Val Leu Ile Asp Ile Val Thr Asp Pro Asn Ala Leu Ser Ile
 525 530 535
 5 cca cca acc atc acg tgg gaa cag gtc atg gga ttc agc aag gcg gcc 1985
 Pro Pro Thr Ile Thr Trp Glu Gln Val Met Gly Phe Ser Lys Ala Ala
 540 545 550
 10 acc cga acc gtc ttt ggt gga gga gta gga gcg atg atc gat ctg gcc 2033
 Thr Arg Thr Val Phe Gly Gly Gly Val Gly Ala Met Ile Asp Leu Ala
 555 560 565
 15 cgt tcg aac ata agg aat att cct act cca tgatgattga tacacctgct 2083
 Arg Ser Asn Ile Arg Asn Ile Pro Thr Pro
 570 575
 20 gttctcattg accgcgagcg cttaactgcc aacatttcca ggatggcagc tcacgccggt 2143
 gcccatgaga ttgccct 2160
 <210> 5
 <211> 579
 <212> PRT
 <213> Corynebacterium glutamicum
 25 <400> 5
 Met Ala His Ser Tyr Ala Glu Gln Leu Ile Asp Thr Leu Glu Ala Gln
 1 5 10 15
 30 Gly Val Lys Arg Ile Tyr Gly Leu Val Gly Asp Ser Leu Asn Pro Ile
 20 25 30
 Val Asp Ala Val Arg Gln Ser Asp Ile Glu Trp Val His Val Arg Asn
 35 35 40 45
 Glu Glu Ala Ala Ala Phe Ala Ala Gly Ala Glu Ser Leu Ile Thr Gly
 50 55 60
 40 Glu Leu Ala Val Cys Ala Ala Ser Cys Gly Pro Gly Asn Thr His Leu
 65 70 75 80
 Ile Gln Gly Leu Tyr Asp Ser His Arg Asn Gly Ala Lys Val Leu Ala
 85 90 95
 45 Ile Ala Ser His Ile Pro Ser Ala Gln Ile Gly Ser Thr Phe Phe Gln
 100 105 110
 Glu Thr His Pro Glu Ile Leu Phe Lys Glu Cys Ser Gly Tyr Cys Glu
 115 120 125
 50 Met Val Asn Gly Gly Glu Gln Gly Glu Arg Ile Leu His His Ala Ile
 130 135 140
 55 Gln Ser Thr Met Ala Gly Lys Gly Val Ser Val Val Val Ile Pro Gly
 145 150 155 160
 Asp Ile Ala Lys Glu Asp Ala Gly Asp Gly Thr Tyr Ser Asn Ser Thr
 165 170 175
 60 Ile Ser Ser Gly Thr Pro Val Val Phe Pro Asp Pro Thr Glu Ala Ala
 180 185 190
 Ala Leu Val Glu Ala Ile Asn Asn Ala Lys Ser Val Thr Leu Phe Cys

	195					200					205					
	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	Gln	Val	Leu	Glu	Leu	Ala	Glu
	210						215					220				
5	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	Leu	Gly	Gly	Lys	Gln	Tyr	Ile
	225					230					235					240
	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	Met	Ser	Gly	Leu	Leu	Gly	Tyr
10					245					250					255	
	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	Ala	Asp	Leu	Leu	Ile	Leu	Leu
				260					265					270		
15	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	Leu	Pro	Lys	Asp	Asn	Val	Ala
			275					280					285			
	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	Gly	Arg	Arg	Thr	Thr	Val	Lys
20		290					295					300				
	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	Thr	Ile	Glu	Asn	Ile	Leu	Pro
	305					310					315					320
	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	Phe	Leu	Asp	Arg	Met	Leu	Lys
25					325					330					335	
	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	Val	Glu	Thr	Tyr	Thr	His	Asn
				340					345					350		
30	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	Glu	Tyr	Val	Ala	Ser	Ile	Leu
			355					360					365			
	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	Phe	Thr	Val	Asp	Thr	Gly	Met
35		370					375					380				
	Cys	Asn	Val	Trp	His	Ala	Arg	Tyr	Ile	Glu	Asn	Pro	Glu	Gly	Thr	Arg
	385					390					395					400
	Asp	Phe	Val	Gly	Ser	Phe	Arg	His	Gly	Thr	Met	Ala	Asn	Ala	Leu	Pro
40					405					410					415	
	His	Ala	Ile	Gly	Ala	Gln	Ser	Val	Asp	Arg	Asn	Arg	Gln	Val	Ile	Ala
				420					425					430		
45	Met	Cys	Gly	Asp	Gly	Gly	Leu	Gly	Met	Leu	Leu	Gly	Glu	Leu	Leu	Thr
			435					440					445			
	Val	Lys	Leu	His	Gln	Leu	Pro	Leu	Lys	Ala	Val	Val	Phe	Asn	Asn	Ser
50		450					455					460				
	Ser	Leu	Gly	Met	Val	Lys	Leu	Glu	Met	Leu	Val	Glu	Gly	Gln	Pro	Glu
	465					470					475					480
	Phe	Gly	Thr	Asp	His	Glu	Glu	Val	Asn	Phe	Ala	Glu	Ile	Ala	Ala	Ala
55					485					490					495	
	Ala	Gly	Ile	Lys	Ser	Val	Arg	Ile	Thr	Asp	Pro	Lys	Lys	Val	Arg	Glu
				500					505					510		
60	Gln	Leu	Ala	Glu	Ala	Leu	Ala	Tyr	Pro	Gly	Pro	Val	Leu	Ile	Asp	Ile
			515					520					525			
	Val	Thr	Asp	Pro	Asn	Ala	Leu	Ser	Ile	Pro	Pro	Thr	Ile	Thr	Trp	Glu

530 535 540
 Gln Val Met Gly Phe Ser Lys Ala Ala Thr Arg Thr Val Phe Gly Gly
 545 550 555 560
 5 Gly Val Gly Ala Met Ile Asp Leu Ala Arg Ser Asn Ile Arg Asn Ile
 565 570 575
 10 Pro Thr Pro
 15 <210> 6
 <211> 875
 <212> DNA
 <213> Corynebacterium glutamicum
 20 <400> 6
 tgcgagatgg tgaatggtgg tgagcagggt gaacgcattt tgcatacacgc gattcagttcc 60
 accatggcgg gtaaagggtgt gtcggtggtg gtgattcctg gtgatatcgc taaggaagac 120
 gcagggtgacg gtacttattc caattccact atttcttctg gcactcctgt ggtgttcccg 180
 gatcctactg aggctgcagc gctggtggag gcgattaaca acgctaagtc tgtcactttg 240
 25 ttctgcggtg cgggcgtgaa gaatgctcgc gcgcagggtg tggagttggc ggagaagatt 300
 aaatcaccca tcgggcatgc gctgggtggt aagcagtaca tccagcatga gaatccgttt 360
 gaggtcggca tgtctggcct gcttgggttac ggcgctgcg tggatgcgtc caatgaggcg 420
 gatctgctga ttctattggg tacggatttc ccttattctg atttccttcc taaagacaac 480
 gttgcccagg tggatatcaa cggcgcgac attggtcgac gtaccacggt gaagtatccg 540
 30 gtgaccggtg atgttgctgc aacaatcgaa aatattttgc ctcatgtgaa ggaaaaaaca 600
 gatcgttcct tccttgatcg gatgctcaag gcacacgagc gtaagttgag ctcggtggta 660
 gagacgtaca cacataacgt cgagaagcat gtgcctattc accctgaata cgttgcctct 720
 attttgaacg agctggcgga taaggatgcg gtgtttactg tggataccgg catgtgcaat 780
 gtgtggcatg cgaggtacat cgagaatccg gagggaaacg gcgactttgt gggttcattc 840
 35 cgccacggca cgatggctaa tgcgttgctt catgc 875